

***In vivo* and *in vitro* parameters for acaricide efficacy of macrocyclic lactones against *Psoroptes ovis* in cattle**

Author

Wouter van Mol¹, Nathalie De Wilde¹, Stijn Casaert¹, Zhen Zhen Chen¹, Marieke Vanhecke¹, Luc Duchateau², Edwin Claerebout¹

Organisation

¹Department of Virology, Parasitology and Immunology, Faculty of Veterinary Medicine, Ghent University

²Department of Nutrition, Genetics and Ethology, Faculty of Veterinary Medicine, Ghent University

Abstract

Psoroptic mange is an important disease in beef cattle, and Belgian Blue cattle are particularly susceptible. Treatment failure of macrocyclic lactones (ML) against *Psoroptes ovis* has been reported, but clear evidence for resistance of *P. ovis* against ML in cattle is lacking. This study was conducted to investigate ML efficacy in 16 beef farms in Belgium and The Netherlands *in vivo* and *in vitro*.

On each farm a group of animals (n= 7-14) with clinical psoroptic mange was treated with two subcutaneous injections of a short-acting ML with 7-10 days interval (n=15) or a single injection with a long-acting ML (n=1). *In vivo* efficacy was assessed by the reduction in mite counts and the cure rate after the first treatment round and the number of treatment rounds needed to cure all animals. *In vitro* knock-down and mortality was evaluated in a contact assay based on Brimer et al., 1995 (Vet Parasitol 59, 249-255).

All farms needed ≥ 2 treatment rounds (2-8) to obtain full efficacy. Cure rates varied from 0%-80%. Only three farms had a mite count reduction of $>90\%$, two of which had a lower limit of the confidence interval of $<90\%$. All other farms had a mite count reduction $<90\%$ (-411%-81%). LD₅₀ values *in vitro* varied from 2,951-36,867 $\mu\text{g/mL}$ and 0.3-58.3 $\mu\text{g/mL}$ at 24h and 120h, respectively. No significant correlation was found between *in vitro* LD₅₀ values and any of the parameters for *in vivo* efficacy.

In conclusion, unambiguous treatment failure was detected on 13/16 beef farms, confirming the presence of ML resistance in Belgian Blue beef farms. Tentative *in vitro* parameters could not detect ML resistance. The potential use of different *in vitro* and *in vivo* parameters to evaluate acaricide efficacy and to detect acaricide resistance will be discussed.